


5H
1
U511
Fishes

INVESTIGATIONS IN FISH CONTROL

96. Effects of Environmental Factors on the Toxicity of Chloramine-T to Fish
97. Effects of Organic Matter and Loading Rates of Fish on the Toxicity of Chloramine-T



UNITED STATES DEPARTMENT OF THE INTERIOR
FISH AND WILDLIFE SERVICE

Investigations in Fish Control, published by the Fish and Wildlife Service, include reports on the results of work at the National Fisheries Research Center at LaCrosse, Wisconsin, and reports of other studies related to that work. Though each report is regarded as a separate publication, several may be issued under a single cover, for economy. See inside back cover for list of current issues.

Copies of this publication may be obtained from the Publications Unit, U.S. Fish and Wildlife Service, Washington, DC 20240, or may be purchased from the National Technical Information Service (NTIS), 5285 Port Royal Road, Springfield, VA 22161.

Library of Congress Cataloging in Publication Data

Effects of environmental factors on the toxicity of chloramine-T to fish.

(*Investigations in fish control*; 96-97)

Supt. of Docs. no.: I 49.70:96-97

1. Chloramine-T—Toxicology. 2. Fishes—Effect of drugs on.
3. Chloramine-T—Dose-response relationship. 4. Chloramine-T—Environmental aspects. I. Bills, T. D. II. Effects of organic matter and loading rates of fish on the toxicity of chloramine-T. 1988. III. Series.

SH177.C39E44 1988 597.024 88-600475


INVESTIGATIONS IN FISH CONTROL

96. Effects of Environmental Factors on the Toxicity of Chloramine-T to Fish

By T. D. Bills, L. L. Marking, V. K. Dawson, and J. J. Rach

97. Effects of Organic Matter and Loading Rates of Fish on the Toxicity of Chloramine-T

By T. D. Bills, L. L. Marking, V. K. Dawson, and G. E. Howe



U.S. Fish and Wildlife Service
National Fisheries Research Center—LaCrosse
P.O. Box 818
LaCrosse, Wisconsin 54602

...of the ...
...of the ...
...of the ...

The ... of ...
... of ...

... of the ...
... of the ...
... of the ...
... of the ...
... of the ...

... of the ...
... of the ...

... of the ...
... of the ...
... of the ...
... of the ...
... of the ...

... of the ...
... of the ...
... of the ...

Effects of Environmental Factors on the Toxicity of Chloramine-T to Fish

by

T. D. Bills, L. L. Marking, V. K. Dawson,
and J. J. Rach

*U.S. Fish and Wildlife Service
National Fisheries Research Center
P.O. Box 818
LaCrosse, Wisconsin 54602-0818*

Abstract

The toxicity of chloramine-T (*n*-sodium-*n*-chloro-para-toluenesulfonamide), a therapeutic used to treat bacterial gill disease in fish, was evaluated under a variety of physical and chemical conditions. The toxicity (96-h LC₅₀) was 2.80 mg/L for rainbow trout, *Salmo gairdneri*; 3.75 mg/L for channel catfish, *Ictalurus punctatus*; and 7.30 mg/L for fathead minnows, *Pimephales promelas*. Chloramine-T was more toxic in warm water than in cold water at exposures of 24 h or less, but temperature had no significant effect on toxicity in 96-h exposures. The chemical was slightly less toxic to rainbow trout in hard than in soft water, but water hardness had little influence on its toxicity to channel catfish. The pH was the most important factor affecting chloramine-T toxicity; the chemical was about 6 times more toxic at pH 6.5 than at 9.5. At levels $\times 1$, $\times 3$, and $\times 5$ the recommended use concentration of 12 mg/L for 1 h, there was no significant mortality in rainbow trout and no abnormal responses were observed in treated fish. In rainbow trout, the toxicity of solutions of chloramine-T aged for 4 weeks was about half that of fresh solutions (deactivation index = 2.10).

Introduction

Chloramine-T (*n*-sodium-*n*-chloro-para-toluenesulfonamide) was shown by From (1980) to be effective for controlling bacterial gill disease (BGD), which is one of the most common diseases of hatchery-reared salmonids and causes more fish losses than any other bacterial disease. The disease is strongly mediated by the stressful environmental conditions and marginal nutrition commonly associated with intensive culture. Flavobacteria and flexibacteria are associated agents (Snieszko 1981). Gills damaged by BGD pathogens are also prone to secondary invasion by fungi (Warren 1981). Because BGD is a limiting factor in fish production, a control agent, such as chloramine-T, is needed for use on food fish. Approval by the U.S. Food and Drug Administration will be required.

Researchers have reported that the activity of chloramine-T is affected by pH, water hardness, and temperature. Tooby et al. (1975) found that the toxicity of chloramine-T to fish and eggs increased as pH and water hardness decreased. Cross and Hursey (1973) observed that chloramine-T was more toxic to fish in soft, acid waters than in hard, alkaline waters, and less toxic at 10° C than at high temperatures.

The purpose of the present study was to delineate the five factors that affect the toxicity of chloramine-T to selected species of fish: (1) sensitivity differences between selected coldwater and warmwater fish species; (2) effects of pH, water hardness, and temperature; (3) persistence of toxicity in solutions over a 4-week period; (4) toxicity and observed effects at recommended use pattern concentrations for rainbow trout (*Salmo gairdneri*); and (5) the time-independent toxicity.

Materials and Methods

Static and flow-through test procedures used in this study followed those prescribed by the Committee on Methods for Toxicity Tests with Aquatic Organisms (1975), ASTM Committee E-35 on Pesticides (1980), and U.S. Department of Agriculture (1986). We exposed 20 fish to each concentration of chloramine-T in glass jars containing 15 L of oxygen-saturated test water. Reconstituted test waters were prepared according to standardized procedures to produce the desired water quality. The pH of test waters was controlled with chemical buffers (Committee on Methods for Toxicity Tests with Aquatic Organisms 1975). The solutions were adjusted to the desired pH before each test began and readjusted with chemical buffers at 24-h intervals, as needed, to maintain the selected pH (± 0.2 unit). Temperatures were regulated by immersing the test jars in constant temperature water baths. To assess the effects of water hardness, we buffered solutions to a constant pH with sodium bicarbonate using the procedure of Marking (1975).

Rainbow trout, fathead minnows (*Pimephales promelas*), and channel catfish (*Ictalurus punctatus*), obtained from a Federal fish hatchery or produced at the National Fisheries Research Center, LaCrosse, Wisconsin, were maintained according to the standard procedures for handling bioassay fish described by Hunn et al. (1968). The fish were acclimated to the desired water chemistries and temperatures for 24 h before each test. Mortalities were recorded at 1, 3, 6, 12, 24, 48, 72, and 96 h.

Two species (rainbow trout and channel catfish) were used in tests to determine the effects of water temperature, hardness, and pH on the toxicity of chloramine-T. In tests on the effects of use pattern levels, we exposed rainbow trout to chloramine-T (12 mg/L for 1 h) and to $\times 3$ and $\times 5$ levels, as prescribed in the IR-4 Guidelines for Investigation of Minor Use Drugs (U.S. Department of Agriculture 1986). We observed responses of fish after exposure for 14 days, using the criteria of Lennon and Walker (1964).

Commercial grade chloramine-T (lot 12605), obtained from Badger Pharmacal Inc. (Jackson, Wisconsin), was accompanied by a certificate of analysis that listed the assay as 100.03% available chlorine. When we tested the material, using the method provided by the manufacturer (U.S. Pharmacopeial Convention, Inc. 1979), the material yielded 98.4% available chlorine.

Concentrations of chloramine-T in test waters were determined by high performance liquid chromatography (HPLC) at 0, 6, 24, 48, 72, and 96 h. Samples were

filtered through 0.45 μ m Acrodiscs, and then injected directly onto the column with an automatic Waters Intelligent Sample Processor (WISP). Retention time was about 3.2 min. Quantification of the peaks was performed by a 730 Data Module having external standard calibration.

The Waters Associates, Inc., high performance liquid chromatography unit that we used consisted of a Model 481 Lambda-Max LC spectrophotometer, Model 510 pump, Model 710B WISP auto sampler, and 730 Data Module. The operating conditions were as follows: stationary phase, 30 cm \times 4 mm Varian MicroPak MCH-10; mobile phase, acetonitrile: phosphate buffer (50:50, v/v); flow rate, 2.0 mL/min; chart speed, 2.0 cm/min; wavelength, 229 nm; and attenuation, 0.10 absorption units. A 0.45- μ m disposal filter assembly was used to filter the sample.

Reagents were acetonitrile, HPLC grade; water, HPLC grade; phosphoric acid, American Chemical Society reagent grade 0.2 M, 13.6 mL diluted to 1 L with HPLC water; monobasic potassium phosphate, American Chemical Society reagent grade 0.2 M, consisting of 27.2 g of KH_2PO_4 diluted to 1 L with HPLC water; and buffer reagent, 0.1 M, consisting of 14.3 mL of 0.2 M H_3PO_4 + 10.7 mL of 0.2 M KH_2PO_4 diluted to 500 mL (pH = 2.6).

In tests to determine the persistence of chloramine-T in water, we aged solutions in glass containers under routine laboratory conditions with 12-h photoperiods for as long as 4 weeks. Residual concentrations of chloramine-T in the aged solutions were determined analytically each week and the toxicity was compared with that of fresh solutions. Deactivation indices were calculated by dividing the 96-h LC50 of the aged solutions by the 96-h LC50 of the fresh solutions (Marking 1972).

The methods of Litchfield and Wilcoxon (1949) were used to compute the LC50's and 95% confidence intervals. Time-independent LC50's (TILC50) were calculated according to the method of Green (1965).

Results and Discussion

Toxicity to Selected Species of Fish

Chloramine-T was toxic to all species exposed in soft water; the 96-h LC50's (mg/L) were 2.80 for rainbow trout, 3.75 for channel catfish, and 7.30 for fathead minnows (Table 1). Rainbow trout (coldwater) and channel catfish (warmwater) were thus twice as sensitive as fathead minnows (warmwater).

Table 3. Toxicity (LC50, mg/L and 95% confidence interval) of chloramine-T to channel catfish in water of selected temperatures, water hardnesses, and pH levels.

Temperature (°C)	Hardness	pH	Duration of test (h)					
			1	3	6	12	24	96
12	Soft	7.5	>60.0	>60.0	>60.0	14.2	10.0	3.75
17	Soft	7.5	>60.0	>60.0	25.3	12.2-16.5	9.07-11.0	3.30-4.26
					21.0-30.4	14.2	7.20	3.73
22	Soft	7.5	>60.0	>60.0	14.0	12.2-16.5	6.78-7.64	3.29-4.22
					12.0-16.3	9.00	5.63	3.80
12	Very soft	8.1	>60.0	>60.0	>60.0	8.51-9.51	5.08-6.24	3.33-4.34
12	Soft	8.1	>60.0	>60.0	>60.0	>60.0	28.0	7.70
							24.1-32.6	6.98-8.49
12	Hard	8.1	>60.0	>60.0	>60.0	>60.0	30.5	11.0
							26.3-35.4	9.13-13.2
12	Very hard	8.1	>60.0	>60.0	>60.0	>60.0	33.0	7.80
							29.2-37.3	6.53-9.31
12	Soft	6.5	>60.0	27.0	10.0	5.60	37.0	9.80
							32.7-41.8	8.64-11.1
12	Soft	8.5	>60.0	>60.0	>60.0	>60.0	2.85	1.75
							2.52-3.22	1.48-2.06
12	Soft	9.5	>60.0	>60.0	>60.0	>60.0	51.5	10.5
							46.8-56.7	9.14-12.1
12	Soft	9.5	>60.0	>60.0	>60.0	>60.0	>60.0	12.3
								9.75-15.5

water (280-320 mg/L as CaCO₃). Tooby et al. (1975) and Cross and Hursey (1973) reported that chloramine-T was more toxic in soft, acid waters than in hard, alkaline waters, but they did not isolate the effects of hardness from those of pH.

The toxicity of chloramine-T increased significantly as the pH of test waters decreased for both species (Tables 2 and 3). At pH 9.5, no rainbow trout or channel catfish died after 24 h of exposure to 60 mg/L, but the chemical was about 6 times more toxic to both species at pH 6.5. The 96-h LC50's for rainbow trout and channel catfish were 1.89 and 1.75 mg/L in acid water (pH 6.5) compared with 10.8 and 12.3 mg/L in alkaline water (pH 9.5), respectively (Tables 2 and 3).

Use Pattern Exposure

Responses of rainbow trout exposed to chloramine-T for 1 h to $\times 1$, $\times 3$, and $\times 5$ the recommended use pattern concentration (12 mg/L) did not differ from those of control fish during the 14-day posttreatment recovery period. These fish were exposed in soft water at 12°C and pH 7.5. Therefore, there was little chance of causing mortality with overtreatment.

Persistence

The toxicity of aged solutions of chloramine-T decreased with aging (Table 4). Toxicity to rainbow trout was decreased by a factor of 1/2 after 4 weeks; the 96-h

Table 4. Toxicity and deactivation index for chloramine-T for rainbow trout in soft water at 12°C.

Aging period (weeks)	96-h LC50 (mg/L) and 95% confidence interval	Deactivation index ^a
0	2.85	1.0
1	2.57-3.16	1.44
	4.10	
2	3.54-4.72	1.33
	3.80	
3	3.34-4.32	2.81
	8.00	
4	6.15-10.4	2.10
	6.00	
	5.27-6.82	

^aLC50 of aged solution
LC50 of fresh solution

Table 5. HPLC analysis of chloramine-T concentrations of 6, 10, and 40 mg/L (calculated) remaining in soft water without fish at 12°C for periods as long as 4 weeks.

Age of solution at sampling time (weeks)	Calculated concentration (mg/L)											
	6	10	40	6	10	40	6	10	40	6	10	40
0	6.12	10.26	40.59	5.84	9.88	39.33	5.78	9.58	37.94	5.88	9.87	39.60
1	5.82	9.82	39.60	5.44	9.37	37.46	5.76	9.73	39.06	5.83	9.85	39.75
2	5.43	9.34	37.48	5.62	9.38	39.22	5.89	9.87	40.29			
3	5.41	9.33	38.98	5.58	9.66	41.11						
4	5.50	9.56	40.81									

LC50 was 6.00 mg/L, in comparison with 2.85 mg/L for freshly prepared solutions (deactivation index = 2.10).

In analytical checks of the concentrations of chloramine-T made at weekly intervals throughout the aging experiment, the measured concentrations (Table 5) did not decrease as rapidly as the toxicity of the aged solutions. For example, the 6 mg/L (calculated) concentration was measured at 6.12 mg/L at time 0 and 5.50 mg/L after 4 weeks of aging. For the same aging period, the toxicity decreased by a factor of 2. When calculated exposure concentrations of 6, 10, and 40 mg/L were analyzed by HPLC at the beginning of all tests to verify the accuracy of the chemical additions, the means and standard deviations (mg/L) for the three analyzed concentrations from 40 tests were 5.90 ± 0.17 , 9.89 ± 0.27 , and 38.9 ± 1.39 .

Time-independent Toxicity

Rainbow trout exposed to chloramine-T in flow-through tests were considerably more resistant than those exposed in static tests. In two separate exposures, the 96-h LC50's were 23.0 and 30.0 mg/L in flow-through tests and 7.0 and 4.6 mg/L in static tests (Table 6). We speculate that chlorine is released from chloramine-T more completely under static conditions and believe that this increased release causes the higher toxicity. Mortalities of fish exposed to chloramine-T did not continue beyond 96 h; the time-independent LC50's were similar to the 96-h LC50's in flow-through toxicity tests (Table 6).

References

ASTM Committee E-35 on Pesticides. 1980. Standard practice for conducting acute toxicity tests with fishes, macroinvertebrates, and amphibians, E729-80. Pages 1-25 in Annual book of ASTM standards, Part 46. End use and consumer products. American Society for Testing and Materials, Philadelphia, Pa.

- Committee on Methods for Toxicity Tests with Aquatic Organisms. 1975. Methods for acute toxicity tests with fish, macroinvertebrates, and amphibians. U.S. Environ. Prot. Agency, Ecol. Res. Serv., EPA-660/3-75-009. 61 pp.
- Cross, D. G., and P. A. Hursey. 1973. Chloramine-T for the control of *Ichthyophthirius multifiliis* (Fouquet). J. Fish Biol. 5:789-798.
- From, J. 1980. Chloramine-T for control of bacterial gill disease. Prog. Fish-Cult. 42:85-86.
- Green, R. H. 1965. Estimation of tolerance over an indefinite time period. Ecology 46:887.
- Hunn, J. B., R. A. Schoettger, and E. W. Whealdon. 1968. Observations on handling and maintenance of bioassay fish. Prog. Fish-Cult. 30:164-167.
- Lennon, R. E., and C. R. Walker. 1964. Laboratories and methods for screening fish-control chemicals. U.S. Fish Wildl. Serv., Invest. Fish Control 1, Circ. 185. 15 pp.
- Litchfield, J. T., Jr., and F. Wilcoxon. 1949. A simplified method of evaluating dose-effect experiments. J. Pharmacol. Exp. Ther. 96:99-113.

Table 6. Acute and chronic (time-independent) toxicity of chloramine-T to rainbow trout in well water at 12°C.

Fish source and type of test	LC50's (mg/L) and 95% confidence intervals	
	96 h	TILC50 ^a
Lot 8675		
Static	4.60	—
	4.14-5.11	— —
Flow-through	30.0	24.3
	27.6-32.6	21.5-27.5
Lot 8706		
Static	7.00	—
	6.44-7.61	— —
Flow-through	23.0	23.0
	20.2-26.2	20.1-26.2

^aTime-independent LC50.

- Marking, L. L. 1972. Methods of estimating the half-life of biological activity of toxic chemicals in water. U.S. Fish Wildl. Serv., Invest. Fish Control 46. 9 pp.
- Marking, L. L. 1975. Toxicological protocol for the development of piscicides. Pages 26-31 in P. H. Eschmeyer, ed. Rehabilitation of fish populations with toxicants: A symposium. Am. Fish. Soc. Spec. Publ. 4.
- Snieszko, S. F. 1981. Bacterial gill disease of freshwater fishes. U.S. Fish Wildl. Serv., Fish Dis. Leaflet. 62. 11 pp.
- Tooby, T. E., P. A. Hursey, and J. S. Alabaster. 1975. The acute toxicity of 102 pesticides and miscellaneous substances to fish. Chem. Ind. (21 June 1975):523-526.
- U.S. Department of Agriculture. 1986. Interregional research project no. 4. Guidelines for IR-4 investigations. New Jersey Agricultural Experiment Station, Cook College, Rutgers University, New Brunswick, N.J. 6pp.
- U.S. Pharmacopeial Convention, Inc. 1979. Chloramine-T assay. Pages 1057-1058 in U.S. Pharmacopoeia. 20th Revision. Mack Publishing Co., Easton, Pa.
- Warren, J. W. 1981. Diseases of hatchery fish. U.S. Fish and Wildlife Service, Twin Cities, Minn. 91 pp.

Bills, Terry D., Leif L. Marking, Verdel K. Dawson, and Jeffery J. Rach. 1988. **Effects of environmental factors on the toxicity of chloramine-T to fish.** U.S. Fish Wildl. Serv., Invest. Fish Control 96. 6 pp.

The toxicity of chloramine-T (*n*-sodium-*n*-chloro-para-toluenesulfonamide), a therapeutic used to treat bacterial gill disease in fish, was evaluated under a variety of environmental conditions. The 96-h LC50 (mg/L) was 2.80 for rainbow trout, *Salmo gairdneri*; 3.75 mg/L for channel catfish, *Ictalurus punctatus*; and 7.30 mg/L for fathead minnows, *Pimephales promelas*. Chloramine-T was more toxic in warm water than in cold water only at exposures of 24 h or less; it was about 6 times more toxic at pH 6.5 than at 9.5, but nontoxic at levels $\times 5$ the use pattern level (12 mg/L; 1 h), and detoxified to half the original activity in about 4 weeks.

Key words: Toxicity, chloramine-T, pH, water hardness, temperature.

Bills, Terry D., Leif L. Marking, Verdel K. Dawson, and Jeffery J. Rach. 1988. **Effects of environmental factors on the toxicity of chloramine-T to fish.** U.S. Fish Wildl. Serv., Invest. Fish Control 96. 6 pp.

The toxicity of chloramine-T (*n*-sodium-*n*-chloro-para-toluenesulfonamide), a therapeutic used to treat bacterial gill disease in fish, was evaluated under a variety of environmental conditions. The 96-h LC50 (mg/L) was 2.80 for rainbow trout, *Salmo gairdneri*; 3.75 mg/L for channel catfish, *Ictalurus punctatus*; and 7.30 mg/L for fathead minnows, *Pimephales promelas*. Chloramine-T was more toxic in warm water than in cold water only at exposures of 24 h or less; it was about 6 times more toxic at pH 6.5 than at 9.5, but nontoxic at levels $\times 5$ the use pattern level (12 mg/L; 1 h), and detoxified to half the original activity in about 4 weeks.

Key words: Toxicity, chloramine-T, pH, water hardness, temperature.

Bills, Terry D., Leif L. Marking, Verdel K. Dawson, and Jeffery J. Rach. 1988. **Effects of environmental factors on the toxicity of chloramine-T to fish.** U.S. Fish Wildl. Serv., Invest. Fish Control 96. 6 pp.

The toxicity of chloramine-T (*n*-sodium-*n*-chloro-para-toluenesulfonamide), a therapeutic used to treat bacterial gill disease in fish, was evaluated under a variety of environmental conditions. The 96-h LC50 (mg/L) was 2.80 for rainbow trout, *Salmo gairdneri*; 3.75 mg/L for channel catfish, *Ictalurus punctatus*; and 7.30 mg/L for fathead minnows, *Pimephales promelas*. Chloramine-T was more toxic in warm water than in cold water only at exposures of 24 h or less; it was about 6 times more toxic at pH 6.5 than at 9.5, but nontoxic at levels $\times 5$ the use pattern level (12 mg/L; 1 h), and detoxified to half the original activity in about 4 weeks.

Key words: Toxicity, chloramine-T, pH, water hardness, temperature.

Bills, Terry D., Leif L. Marking, Verdel K. Dawson, and Jeffery J. Rach. 1988. **Effects of environmental factors on the toxicity of chloramine-T to fish.** U.S. Fish Wildl. Serv., Invest. Fish Control 96. 6 pp.

The toxicity of chloramine-T (*n*-sodium-*n*-chloro-para-toluenesulfonamide), a therapeutic used to treat bacterial gill disease in fish, was evaluated under a variety of environmental conditions. The 96-h LC50 (mg/L) was 2.80 for rainbow trout, *Salmo gairdneri*; 3.75 mg/L for channel catfish, *Ictalurus punctatus*; and 7.30 mg/L for fathead minnows, *Pimephales promelas*. Chloramine-T was more toxic in warm water than in cold water only at exposures of 24 h or less; it was about 6 times more toxic at pH 6.5 than at 9.5, but nontoxic at levels $\times 5$ the use pattern level (12 mg/L; 1 h), and detoxified to half the original activity in about 4 weeks.

Key words: Toxicity, chloramine-T, pH, water hardness, temperature.

Effects of Organic Matter and Loading Rates of Fish on the Toxicity of Chloramine-T

by

T. D. Bills, L. L. Marking, V. K. Dawson, and G. E. Howe

*U.S. Fish and Wildlife Service
National Fisheries Research Center
P.O. Box 818
LaCrosse, Wisconsin 54602-0818*

Abstract

Chloramine-T (*n*-sodium-*n*-chloro-para-toluenesulfonamide) is effective for the control of bacterial gill disease in fishes but data on its toxicity and safety are lacking. We examined the effect of fish loading rates, feed levels, and fecal material on the availability and toxicity of the chemical. Its toxicity increased (24-h LC50's were 36 mg/L and 14 mg/L) as loading rates of fathead minnows (*Pimephales promelas*) increased from 0.52 to 2.07 g/L of water. The presence of fish food decreased the toxicity of the chemical. The 96-h LC50 was 7.30 mg/L without feed and 14.2 and 20.8 mg/L after the addition of 0.07 and 0.20 g/L of fish food. The presence of fecal material also significantly decreased the toxicity of chloramine-T. Concentrations in water decreased more rapidly when fish were present than when only feed or fecal matter was present. The decreases were greatest in the presence of fish and feed or fish and fecal material.

Introduction

Chloramine-T (*n*-sodium-*n*-chloro-para-toluenesulfonamide) is an antibacterial agent used as a disinfectant and preservative in the food industry. It is also effective for controlling bacterial gill disease (BGD) in fish (From 1980). Although no single pathogen seems to be responsible for BGD infections, all known agents are gram-negative bacteria—myxobacteria, aeromonads, and pseudomonads (Snieszko 1981). The disease is highly contagious among cultured salmonids and is associated with crowded rearing conditions and sometimes inadequate nutrition. The disease can lead to substantial fish losses. An approved therapeutic to control BGD is needed to enable the production of salmonids for the restoration of fish stocks and for the sport and commercial fisheries.

Several researchers have reported that the activity of chloramine-T is affected by environmental factors. Van Duijn (1967) reported that it is partly or totally inactivated by organic detritus. Similarly, Cross and Hursey

(1973) demonstrated that chloramine-T lost its activity in the presence of excessive organic matter and showed that its toxicity to fish was greater in soft, acid waters than in hard, alkaline waters. Bills et al. (1988) reported that the acute toxicity of chloramine-T was greater at a given concentration in warm than in cold water and that the compound was more toxic in acid than in alkaline water.

The purpose of the present study was to determine (1) the effects of fish loading rates, feed levels, and fecal material on the toxicity of chloramine-T to fish; and (2) the available concentration of chloramine-T in water containing fish, feed, and fecal material.

Materials and Methods

Static test procedures used in this study followed those prescribed by the Committee on Methods for Toxicity with Aquatic Organisms (1975), ASTM Committee E-35 on Pesticides (1980), and U.S. Department of Agriculture

(1986). Reconstituted test water (total hardness, 40–44 mg/L as CaCO_3) was prepared according to standardized procedures. Glass jars containing 15 L of oxygen-saturated water were used for all tests. Exposure temperatures were regulated by immersing the test jars in constant temperature water baths at 12° C. All tests were conducted in duplicate. Twenty fish were exposed to each concentration, except in the fish-loading rate study, in which 7, 14, or 28 fish were used.

Fathead minnows (*Pimephales promelas*) cultured at the National Fisheries Research Center, LaCrosse, Wisconsin, were used as the test organisms. They were maintained according to the standard procedures for handling bioassay fish described by Hunn et al. (1968) and were acclimated to test conditions for 24 h before each test. Mortalities were recorded at 1, 3, 6, 24, 48, 72, and 96 h.

Commercial grade chloramine-T (lot 12605), obtained from Badger Pharmaceutical, Inc. (Jackson, Wisconsin), was used for all tests. The certificate of analysis supplied by the company listed the assay as 100.03% available chlorine. However, the material yielded 98.4% available chlorine when we analyzed it according to the method provided by the manufacturer (U.S. Pharmacopeial Convention, Inc. 1979). All chloramine-T concentrations are expressed in milligrams per liter. Concentrations in selected test containers were checked analytically at 0, 6, 24, 48, 72, and 96 h.

Fathead minnows averaging 1.11 g were used to assess the effects of three loading rates of fish on the toxicity of chloramine-T. We used either 7, 14, or 28 fish to provide loading rates of 0.52, 1.04, or 2.07 g/L. Each set was exposed to a series of concentrations of the test chemical.

We tested the effects of the presence of fish food on the toxicity of chloramine-T by using concentrations of $\times 1$ (0.07 g/L), $\times 2$, and $\times 3$ the recommended feeding rate for fish at 12° C—about 5% of their body weight per day (Piper et al. 1982). On the basis of the weight of the fish, 0.07, 0.13, or 0.20 g/L of Nelsons' Sterling Silver Cup No. 1 fish food was added to each test vessel. The feed contained crude protein 48%, crude fat 14%, crude fiber 3.6%, and ash 12.0%. Chloramine-T was administered immediately after the feed was added.

The effects of fecal matter on the toxicity of chloramine-T were determined by adding three concentrations of a slurry of feces and bottom materials (referred to here as fecal material) collected from a fish production raceway. To determine the amount of fecal material present in the slurry, we drained the excess water and dried five 5-mL portions of the material in an oven at 100° C for 24 h; the mean dry weight per portion was

0.08 g/mL. We then added portions of the wet fecal material to test vessels to yield dry weight equivalent concentrations of 0.017, 0.033, and 0.067 g/L of the material. After adding the material, we manually stirred the contents of each jar, added the fish, and then the chloramine-T at concentrations of 2 to 80 mg/L.

Concentrations of chloramine-T in the test waters were determined by high performance liquid chromatography (HPLC), according to methods described by Bills et al. (1988).

Statistical Analysis

The methods of Litchfield and Wilcoxon (1949) were used to compute the LC_{50} 's (concentration calculated to produce 50% mortality) and 95% confidence intervals.

Results and Discussion

Fish Loading Rates

The toxicity of chloramine-T increased as the fish loading rate increased from 0.52 to 2.07 g/L; 24-h LC_{50} 's were 36.0 and 14.0 mg/L, respectively (Table 1). However, the 96-h LC_{50} 's were not significantly different among the three loading rates. The higher loading rates may have stressed the fish and increased the toxicity, even though oxygen concentrations were maintained above stressful levels (>50% saturation) for fathead minnows. Concentrations of chloramine-T decreased over time—especially at the higher fish loading rates (Table 2). The additional fish apparently absorbed more of the chemical from solution. The decreased chloramine-T concentrations may have offset the increased toxicity associated with crowding. By 96 h, there were no significant differences in toxicity among the different loading rates.

Table 1. Toxicity of chloramine-T to fathead minnows at three different loading rates of fish in soft water at 12° C.

Fish loading rate (g/L)	LC_{50} (mg/L) and 95% confidence interval	
	24 h	96 h
0.52	36.0 31.3–41.3	8.50 7.24–9.98
1.04	22.5 19.6–25.8	7.00 6.52–7.51
2.07	14.0 10.0–19.4	8.00 7.49–8.54

Table 2. Concentrations (mg/L) of chloramine-T remaining after treatments with the chemical in the presence of fish, fish feed, fecal material, fish and fish feed, and fish and fecal material in soft water at 12°C.

Time Time (h)	Treat- ment concen- tration (mg/L)	Fish (g/L)			Fish feed (g/L)			Fecal material (g/L)			Fish and fish feed (g/L)			Fish and fecal material (g/L)				
		0.52	1.04	2.07	0.07	0.13	0.20	0.02	0.03	0.07	Refer- ence ^a	0.07	0.13	0.20	Refer- ence ^a	0.02	0.03	0.07
0	6	5.92	5.93	5.74	4.45	3.37	2.28	5.01	4.89	4.75	5.73	4.46	3.28	2.30	4.99	5.29	5.15	5.04
6	6	5.92	6.54	5.96	4.14	3.23	2.18	5.14	4.81	4.78	5.66	4.31	3.46	2.27	5.15	5.09	4.83	4.68
24	6	5.64	5.34	5.10	3.57	2.46	1.38	5.16	4.68	4.60	5.44	3.43	2.08	1.12	5.22	4.99	4.64	4.41
48	6	5.40	4.94	4.33	2.88	1.51	0	5.05	3.91	4.08	5.05	2.38	1.04	0	4.78	4.40	3.98	3.65
72	6	4.91	4.24	3.56	2.17	0.96	0	4.86	3.29	3.55	4.70	1.64	0	0	4.19	3.91	3.39	3.03
96	6	4.54	3.93	2.94	1.44	1.02	0	4.51	2.81	3.12	4.08	0.91	0	0	3.57	3.35	2.66	2.38
0	10	9.96	10.05	9.70	7.66	5.80	4.53	8.43	8.14	8.09	9.67	7.98	6.08	4.52	8.64	8.95	8.68	8.48
6	10	9.87	9.89	9.35	7.51	5.52	4.29	8.56	8.40	8.02	9.58	7.55	5.83	4.18	8.57	8.44	8.18	7.96
24	10	9.33	9.03	8.10	6.62	4.24	3.04	8.67	8.34	7.81	9.08	6.14	3.98	2.55	8.59	8.29	7.83	7.41
48	10	8.79	7.92	6.54	5.56	2.92	1.71	8.37	7.87	7.14	8.21	4.63	2.38	1.17	7.76	7.43	6.86	6.31
72	10	8.01	— ^b	5.24	4.85	2.03	0.96	8.19	7.47	6.45	7.54	3.57	1.38	0	6.86	6.76	6.28	5.53
96	10	7.74	—	5.61	4.05	1.26	0	7.77	7.02	5.87	—	2.53	0.60	0	6.19	5.55	5.55	4.74
0	40	39.84	38.94	38.57	34.37	30.99	26.42	34.17	34.68	32.31	39.07	34.08	28.31	25.61	35.38	35.26	34.25	34.11
6	40	39.57	38.55	37.50	34.68	30.01	25.34	34.64	34.15	32.06	38.59	32.27	27.09	23.68	34.90	34.24	32.91	32.37
24	40	36.97	— ^b	—	32.96	27.98	23.48	35.81	35.83	33.17	—	—	—	—	—	—	—	—
48	40	—	—	—	31.45	25.14	20.37	35.04	34.98	32.13	—	—	—	—	—	—	—	—
72	40	—	—	—	30.55	24.09	18.17	35.07	34.34	31.43	—	—	—	—	—	—	—	—
96	40	—	—	—	29.27	21.94	15.13	34.57	33.72	30.35	—	—	—	—	—	—	—	—

^aReference solution contained only water and chloramine-T.

^bAll fish died.

Fish Feed

Test solutions with added fish feed were less toxic than those without feed—especially after 96 h of exposure (Table 3): The 96-h LC50's were 7.30 mg/L for solutions without feed, but 20.8 mg/L for solutions containing 0.20 g/L of feed. The presence of fish feed resulted in a loss of chloramine-T. At the 6 mg/L concentration, the addition of 0.07 g/L of fish feed decreased chloramine-T concentrations 80% within 96 h and 0.20 g/L decreased the concentrations to below the detection limit within 48 h. At the 10 mg/L concentration of chloramine-T, the loss was significantly higher when 0.13 or 0.20 g/L of feed was added than when 0.07 g/L was added (Table 2). We believe the loss of chloramine-T from solution was due to adsorption on feed particles as well as to depletion of chlorine due to the oxidation of the organic material.

Table 3. Toxicity of chloramine-T to fathead minnows in soft water containing four different amounts of fish feed.

Amount of feed (g/L)	LC50 (mg/L) and 95% confidence interval	
	24 h	96 h
0	32.3 25.5–40.9	7.30 6.71–7.94
0.07	28.5 24.5–33.1	14.2 12.5–16.1
0.13	28.4 24.4–33.0	17.2 14.4–20.4
0.20	28.0 24.1–32.6	20.8 17.9–24.1

Fecal Material

The presence of fecal material decreased the toxicity of the chloramine-T solutions in 96-h exposures (Table 4). The 96-h LC50's for the solutions containing the three levels of fecal material ranged from 12.0 to 17.4 mg/L, in comparison with 7.40 mg/L for solutions that contained no fecal material. However, there were no significant differences among the LC50's at the 24-h exposures. The presence of fecal material decreased the chloramine-T concentrations at all three loading levels. As fecal material levels increased, the loss of chloramine-T also increased. The decrease in chloramine-T concentration was independent of the initial concentrations and was directly correlated with the amount of fecal material added (Table 2).

Fish Plus Fish Feed

The presence of fish in combination with feed caused the largest loss of chloramine-T observed in any of the combinations of the materials tested (Table 2). The loss of chloramine-T was 100% over 72 h at 6 mg/L with 0.13 g/L feed plus fish, and 100% over 48 h at 0.20 g/L. Reductions were most rapid between 0 and 24 h. The higher the level of feed added, the more rapid the reduction in chloramine-T concentration. Therefore, chloramine-T would be more effective if the fish were not fed before treatment.

Fish Plus Fecal Material

In most tests, the presence of fish in combination with fecal material decreased chloramine-T concentrations to levels similar to those caused by fish plus feed, although the reductions were less significant (Table 2). In 96 h, the 6 mg/L concentration of chloramine-T was decreased by about 67% and the 10 mg/L concentration was reduced by about 50%. In general, fish culturists must be aware that treatments with chloramine-T may not be effective if the system contains high levels of oxidizable material or material that can absorb the chemical and reduce its effective concentration.

References

ASTM Committee E-35 on Pesticides. 1980. Standard practice for conducting acute toxicity tests with fishes, macroinvertebrates, and amphibians, E729-80. Pages 1-25 in *Annual book of ASTM standards*, Part 46. End use and consumer

Table 4. Toxicity of chloramine-T to fathead minnows in soft water containing four different amounts of fish fecal material.

Amount of fecal material g/15 L	LC50 (mg/L) and 95% confidence interval	
	24 h	96 h
0	26.5 20.7-33.9	7.40 6.65-8.23
0.02	25.5 21.1-30.8	12.0 10.3-13.9
0.03	24.5 20.4-29.4	12.2 10.8-13.8
0.07	25.5 21.1-30.8	17.4 14.6-20.7

- products. American Society for Testing and Materials, Philadelphia, Pa.
- Bills, T. D., L. L. Marking, V. K. Dawson, and J. J. Rach. 1988. Effects of environmental factors on toxicity of chloramine-T to fish. U.S. Fish Wildl. Serv., Invest. Fish Control. 96. 6 pp.
- Committee on Methods for Toxicity Tests with Aquatic Organisms. 1975. Methods for acute toxicity tests with fish, macroinvertebrates, and amphibians. U.S. Environ. Prot. Agency, Ecol. Res. Serv., EPA-660/3-75-009. 61 pp.
- Cross, D. G., and P. A. Hursey. 1973. Chloramine-T for the control of *Ichthyophthirius multifiliis* (Fouquet). J. Fish Biol. 5:789-798.
- From, J. 1980. Chloramine-T for control of bacterial gill disease. Prog. Fish-Cult. 42:85-86.
- Hunn, J. B., R. A. Schoetger, and E. W. Whealdon. 1968. Observations on handling and maintenance of bioassay fish. Prog. Fish-Cult. 30:164-167.
- Litchfield, J. R., Jr., and F. Wilcoxon. 1949. A simplified method of evaluating dose-effect experiments. J. Pharmacol. Exp. Ther. 96:99-113.
- Piper, R. G., I. B. McElwain, L. E. Orme, J. P. McCraren, L. G. Fowler, and J. R. Leonard. 1982. Fish hatchery management. U.S. Fish Wildl. Serv., Washington, D.C. 517 pp.
- Snieszko, S. F. 1981. Bacterial gill disease of freshwater fishes. U.S. Fish Wildl. Serv., Fish Dis. Leaflet. 62. 11 pp.
- U.S. Department of Agriculture. 1986. Interregional research project no. 4. Guidelines for IR-4 investigations. New Jersey Agricultural Experiment Station, Cook College, Rutgers University, New Brunswick, N.J. 6 pp.
- U.S. Pharmacopeial Convention, Inc. 1979. Chloramine-T assay. Pages 1057-1058 in U.S. Pharmacopoeia. 20th Revision. Mack Publishing Co., Easton, Pa.
- Van Duijn, C. 1967. Diseases of fishes, 2nd ed. Charles C. Thomas, Publisher, Springfield, Ill. 399 pp.

Bills, Terry D., Leif L. Marking, Verdel K. Dawson, and George E. Howe. 1988. **Effects of organic matter and loading rates of fish on the toxicity of chloramine-T.** U.S. Fish Wildl. Serv., Invest. Fish Control 97. 4 pp.

The present study was designed to determine the effects of different fish loading rates, feed levels, and residual fecal material on availability and toxicity of chloramine-T (*n*-sodium-*n*-chloro-para-toluenesulfonamide), a chemical used for the control of bacterial gill disease. Toxicity of chloramine-T increased as loading rates of fathead minnows (*Pimephales promelas*) increased from 0.52 to 2.07 g/L of water in 24-h exposures. The 96-h LC50's were 7.30 mg/L for solutions without feed and 14.2 to 20.8 mg/L after the addition of 0.07 to 0.20 g/L of feed. The concentration of chloramine-T in water decreased more rapidly in the presence of fish than in the presence of only feed or fecal material. The losses of chloramine-T were greatest in the presence of combinations of fish and feed or fish and fecal material.

Key words: Toxicity, chloramine-T, fish, inactivation, organic matter.

Bills, Terry D., Leif L. Marking, Verdel K. Dawson, and George E. Howe. 1988. **Effects of organic matter and loading rates of fish on the toxicity of chloramine-T.** U.S. Fish Wildl. Serv., Invest. Fish Control 97. 4 pp.

The present study was designed to determine the effects of different fish loading rates, feed levels, and residual fecal material on availability and toxicity of chloramine-T (*n*-sodium-*n*-chloro-para-toluenesulfonamide), a chemical used for the control of bacterial gill disease. Toxicity of chloramine-T increased as loading rates of fathead minnows (*Pimephales promelas*) increased from 0.52 to 2.07 g/L of water in 24-h exposures. The 96-h LC50's were 7.30 mg/L for solutions without feed and 14.2 to 20.8 mg/L after the addition of 0.07 to 0.20 g/L of feed. The concentration of chloramine-T in water decreased more rapidly in the presence of fish than in the presence of only feed or fecal material. The losses of chloramine-T were greatest in the presence of combinations of fish and feed or fish and fecal material.

Key words: Toxicity, chloramine-T, fish, inactivation, organic matter.

Bills, Terry D., Leif L. Marking, Verdel K. Dawson, and George E. Howe. 1988. **Effects of organic matter and loading rates of fish on the toxicity of chloramine-T.** U.S. Fish Wildl. Serv., Invest. Fish Control 97. 4 pp.

The present study was designed to determine the effects of different fish loading rates, feed levels, and residual fecal material on availability and toxicity of chloramine-T (*n*-sodium-*n*-chloro-para-toluenesulfonamide), a chemical used for the control of bacterial gill disease. Toxicity of chloramine-T increased as loading rates of fathead minnows (*Pimephales promelas*) increased from 0.52 to 2.07 g/L of water in 24-h exposures. The 96-h LC50's were 7.30 mg/L for solutions without feed and 14.2 to 20.8 mg/L after the addition of 0.07 to 0.20 g/L of feed. The concentration of chloramine-T in water decreased more rapidly in the presence of fish than in the presence of only feed or fecal material. The losses of chloramine-T were greatest in the presence of combinations of fish and feed or fish and fecal material.

Key words: Toxicity, chloramine-T, fish, inactivation, organic matter.

Bills, Terry D., Leif L. Marking, Verdel K. Dawson, and George E. Howe. 1988. **Effects of organic matter and loading rates of fish on the toxicity of chloramine-T.** U.S. Fish Wildl. Serv., Invest. Fish Control 97. 4 pp.

The present study was designed to determine the effects of different fish loading rates, feed levels, and residual fecal material on availability and toxicity of chloramine-T (*n*-sodium-*n*-chloro-para-toluenesulfonamide), a chemical used for the control of bacterial gill disease. Toxicity of chloramine-T increased as loading rates of fathead minnows (*Pimephales promelas*) increased from 0.52 to 2.07 g/L of water in 24-h exposures. The 96-h LC50's were 7.30 mg/L for solutions without feed and 14.2 to 20.8 mg/L after the addition of 0.07 to 0.20 g/L of feed. The concentration of chloramine-T in water decreased more rapidly in the presence of fish than in the presence of only feed or fecal material. The losses of chloramine-T were greatest in the presence of combinations of fish and feed or fish and fecal material.

Key words: Toxicity, chloramine-T, fish, inactivation, organic matter.

(Reports 87 through 89 are in one cover.)

87. Ethyl-*p*-aminobenzoate (Benzocaine): Efficacy as an Anesthetic for Five Species of Freshwater Fish, by V. K. Dawson and P. A. Gilderhus. 1979. 5 pp.
88. Influences of Selected Environmental Factors on the Activity of a Prospective Fish Toxicant, 2-(Digeranyl-amino)-ethanol, in Laboratory Tests, by C. A. Launer and T. D. Bills. 1979. 4 pp.
89. Toxicities of the Lampricides 3-Trifluoromethyl-4-nitrophenol (TFM) and the 2-Aminoethanol Salt of 2',5-Dichloro-4'-nitrosalicylanilide (Bayer 73) to Four Bird Species, by R. H. Hudson. 1979. 5 pp.

(Reports 90 and 91 are in one cover.)

90. Accumulation and Loss of 2',5-Dichloro-4'-nitrosalicylanilide (Bayer 73) by Fish: Laboratory Studies, by V. K. Dawson, J. B. Sills, and Charles W. Luhning. 1982. 5 pp.
91. Effects of Synergized Rotenone on Nontarget Organisms in Ponds, by R. M. Burress. 1982. 7 pp.

(Reports 92 through 94 are in one cover.)

92. Acute and Chronic Toxicity of Rotenone to *Daphnia magna*, by J. J. Rach, T. D. Bills, and L. L. Marking. 1988. 5 pp.
93. Toxicity of Rotenone to Developing Rainbow Trout, by T. D. Bills, J. J. Rach, and L. L. Marking. 1988. 3 pp.
94. Oral Toxicity of Rotenone to Mammals, by L. L. Marking. 1988. 5 pp.
95. Deposition and Persistence of Rotenone in Shallow Ponds During Cold and Warm Seasons, by P. A. Gilderhus, V. K. Dawson, and J. L. Allen. 1988. 7 pp.



TAKE PRIDE

in America



U.S. DEPARTMENT OF THE INTERIOR
FISH AND WILDLIFE SERVICE



As the Nation's principal conservation agency, the Department of the Interior has responsibility for most of our nationally owned public lands and natural resources. This includes fostering the wisest use of our land and water resources, protecting our fish and wildlife, preserving the environmental and cultural values of our national parks and historical places, and providing for the enjoyment of life through outdoor recreation. The Department assesses our energy and mineral resources and works to assure that their development is in the best interests of all our people. The Department also has a major responsibility for American Indian reservation communities and for people who live in island territories under U.S. administration.